

[Web](#) [Images](#) [Videos](#) [Maps](#) [News](#) [Shopping](#) [Gmail](#) [more ▼](#)
[Sign in](#)

Google scholar

[Advanced Scholar Search](#)
[Scholar Preferences](#)
ScholarResults **41 - 44** of **44**. (0.14 sec)

Computer programs for calculation of **median effective dose** (LD 50 or ED 50) using the ...

MM Schaper, RD Thompson, CS Weil - Archives of toxicology, 1994 - Springer
 Abstract. Over the past 40 years, toxicologists and pharmacologists have used tables published by Weil for the determination of LDs₀ (or EDs₀) values and their associated 95% confidence intervals. With the advances in computer ...

[Cited by 2](#) - [Related articles](#) - [BL Direct](#) - [All 3 versions](#)

The **median effective dose** of nasal midazolam for endoscopic examination in pediatric ...

ES Moran, GM Chavira, FG Butron Lopez, CM ... - Current Therapeutic Research, 1995 - Elsevier
 Midazolam, a benzodiazepine, was recently included in the basic drug scheme of the Mexican health system. Midazolam shares anxiolytic, sedative, hypnotic, muscle relaxant, amnesic, and anticonvulsant activity with other ...

[Related articles](#) - [BL Direct](#) - [All 3 versions](#)

[CITATION] Computer programs for calculation of **median effective dose**(LD

MM Schaper, RD Thompson, CS Weil - Archives of Toxicology, 1994

Median effective dose of propofol required for induction of anesthesia in goats.

PLSBJEK JCH - Journal of the American Veterinary Medical Association, 1997 - 万方数据资源系统

The **median effective dose** (ED₅₀) of propofol required for induction of anaesthesia and the frequency of myoclonic activity and apnoea associated with propofol administration was determined using 28 goats. The first goat was ...



 Result Page: [Previous](#) [1](#) [2](#) [3](#) [4](#) [5](#)

[Go to Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2009 Google